



Single Top Quark Physics at the Tevatron and Beyond

Catalin Ciobanu, University of Illinois, CDF

For the LHC part, valuable help received from:

Sergey Sablospitsky, IHEP Protvino, CMS

Dugan O'Neil, Simon Fraser Univ., ATLAS

WIN'03, October 7, 2003



Single Top Production

Why look for single top:


- Cross section $\sim |V_{tb}|^2$: the only way to directly measure CKM matrix el. V_{tb}
- Single top is background to other signals: e.g. Higgs searches.
- Test non-SM phenomena:
 - Heavy W' boson
 - Anomalous Wtb couplings
 - FCNC couplings tug or tcb

At the Tevatron relevant channels are:

- t-channel W-gluon fusion:
 - 1.98 pb at $\sqrt{s}=1.96$ TeV
 - Hard b-jet, W decay products, soft b (usually lost), light quark jet
- s-channel W^* :
 - 0.88 pb at $\sqrt{s}=1.96$ TeV
 - Two hard b-jets, W decay products


➤ [B.W. Harris et al.: Phys. Rev. D 66, 054024 \(theoretical calculations\)](#)

W-gluon fusion process



Signal is $W+b+q$

s-channel W^ process*



Signal is $W+b+b$



From Run I to Run II

- Results at $\sqrt{s} = 1.8$ TeV from CDF and DØ :
 - Single top has not been observed; 95% CL limits were set:
 - t-channel: DØ limit: **22 pb**, CDF limit: **13 pb** (theoretical x-section: **1.40 pb**)
 - s-channel: DØ limit: **17 pb**, CDF limit: **18 pb** (theoretical x-section: **0.76 pb**)
 - combined s- and t- channels search: CDF limits: 14 pb (H_T , M_{lnb})
 - most recent Run 1 CDF study finds a 2.2σ excess and limit of 24 pb (7-inp NN)
 - DØ: Phys. Lett. B 517, 282 (2001); CDF: PRD65, 091102 (2002), C. Ciobanu's Ph.D. thesis
- For Run II:
 - Higher rate: 32% increase in the combined s and t channel x-section:
 - still small: 2.9 pb versus 6.7 pb for top pair production
 - More luminosity – right now CDF has 180 pb^{-1} – that is 70% more data!
 - Better detector acceptance
 - Preliminary CDF analysis done for 107 pb^{-1} of data:
 - Search for s- and t-channels combined production
 - Seaparate search for t-channel



Combined Search

- Look in the W+2 tight jets channel:
 - “tight” jet: $E_T > 15$ GeV, $|\eta| < 2.8$
 - at least one SVX B-tag
 - exactly one lepton with $E_T > 20$ GeV, $|\eta| < 2.0$
 - missing energy: $\cancel{E}_T > 20$ GeV
 - veto Z's, dilepton events
 - apply a top mass window cut: $140 < M_{\text{Inb}} < 210$ GeV/c²
- Monte Carlo samples:
 - single top signal:
 - Pythia (1M events)
 - Madevent+Pythia (~200k)
 - HERWIG tt background
 - ALPGEN+HERWIG Wbb used to model the shapes of the non-top backgrounds: Wbb, Wcc, Wc, mistags, non-W events



Expected Yield



- For $L=107.1 \text{ pb}^{-1}$:
 - expect 2.4 signal events
 - expect 16.1 backgrd events
 - observe 19 (expect 18.5)
- The variable with the most discrimination power is H_T , the total transverse energy in the event:
 - Will fit the H_T shape in data as the sum of weighted MC shapes.
 - Use maximum likelihood method with Gaussian background constraints

Process	N in 107.1 pb ⁻¹
W-gluon	1.61 ± 0.51
W*	0.80 ± 0.18
tt	2.33 ± 0.70
Wbb	3.5 ± 1.5
Wcc	1.4 ± 0.8
Wc	3.1 ± 1.0
Mistags	2.9 ± 0.9
Non-W	2.2 ± 0.7
Diboson	0.7 ± 0.2
Z+bb	0.07 ± 0.03
Total non-top	13.8 ± 2.8



Fitting H_T distributions

- H_T distributions for signal and backgrounds:
 - signal shapes similar for s- and t- channels
 - fit yields 2.9 ± 4.5 signal events (exp. 2.4 events).






95% C.L. limit

- Limit at 95% C.L. is 17.5 pb
- This accounts for the systematics:
 - Jet E_T scale: 20.9%
 - ISR/FSR: 4.0%
 - Top Mass: 6.2%
 - PDF, signal generator, background model 1-2%
- Compare to Run I:
 - Same luminosity 107 vs 106 (pb^{-1})
 - 14 pb limit for 2.16 pb x-section
 - Now: 17.5 pb limit for 2.86 pb x-s.

marginally better limit!

- Several ways to improve this will be discussed later.







Separate Search

- Looking for the two signal channels individually is desired:
 - Different sensitivities to new physics:
 - s-channel: heavy charged vector bosons W' , CP-violation effects within MSSM, Kaluza-Klein excited W-boson within MSSM
 - t-channel: FCNC couplings, anomalous V+A contributions to the W-t-b vertex, etc.
 - To go back to extracting V_{tb} we will have to know the individual rates
- We use the same selection as in the combined search.
 - However, using H_T is no longer appropriate
 - A variable with good W^* - W-gluon separation potential is $Q \times \eta$, where Q is the lepton charge and η the pseudorapidity of the non-b jet.
 - Now treat W^* as a background, therefore perform a 4-component fit



Fitting $Q \times \eta$

- t-channel exhibits an asymmetry toward $Q \times \eta$ positive values (twice as many events than in the negative $Q \times \eta$ range)
- fit to data yields: 1.9 ± 3.7 t-channel signal events (exp. 1.6 events).






95% C.L. limit

- Limit at 95% C.L. is 15.4 pb
- This accounts for the systematics:
 - Jet E_T scale: 21.5%
 - Signal generator: 21.1%
 - Background model: 43.0%
 - ISR/FSR: 6.9%
 - Top Mass: 6.7%
 - PDF 4.3%
- Compare to Run I:
 - 13 pb limit for 1.40 pb x-section
 - Now: 15.4 pb limit for 1.98 pb x-s.


again a better limit!





Future Prospects

- In progress: Neural Network with 5 input variables: H_T , $E_T(j1)$, $E_T(j2)$, E_T , $P_T(j1-j2)$, Looking in the $W+2j$ channel, no $M_{l\nu b}$ cut
Assuming the background uncertainties will scale as $L^{-1/2}$




- Right now: 3 sigma with 2.5 fb⁻¹
- Improvements to come. We try:
 - Soft lepton tagger, Jet Prob. tagger
 - Use of forward electrons
 - Better discriminant variables:
 - Matrix elements (or ratios of these)
 - Use the top mass hypothesis to constrain the event
 - Will lead to better $M_{l\nu b}$ resolution
 - Better understanding of the systematic uncertainties is crucial.



Single Top at the LHC


- LHC will be a top quark factory:
 - 8 million top pairs per experiment per year ($10 \text{ fb}^{-1} / \text{year}$)
 - Some cross section values for $\sqrt{s} = 14 \text{ TeV}$:
 - top pair production: $\sim 800 \text{ pb}$ (mostly via gluon-gluon fusion)
 - t-channel single top: 153 (top) and 90 (antitop) = 243 pb
 - **e.g. per day ~ 6000 events, at $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$**
 - s-channel single top: 6.6 (top) and 4.8 (antitop) = 11 pb
 - associated Wt production: 50-60 pb
 - **Negligible at the Tevatron**
- Will discuss each of the three modes in what follows





t-channel single top (CMS)

- CMS study (hep-ph/0003033):
 - Full calorimeter simulation + b-tag efficiency parametrization
 - Pythia 5.72 signal, tt, WZ; Vecbos+Herwig W+jets. Selection:
 - One isolated lepton with $P_T > 20$ GeV/c, $|\eta| < 2.5$
 - Missing $E_T > 20$ GeV, $50 < M_{l\nu} < 100$ GeV/c²
 - Two jets with $E_T > 20$ GeV, $|\eta| < 4.0$
 - **One jet: $E_T > 20$ GeV, $|\eta| < 2.5$, the other $E_T > 50$ GeV, $2.5 < |\eta| < 4.0$**
 - **Leading jet $E_T < 100$ GeV (to reduce tt)**
 - **Exactly one b-tagged jet – the central one**
 - Reject WZ candidates with: $80 < M_{jj} < 100$ GeV/c²
 - Signal peak visible in reconstructed M_{top} distribution
 - With 10 fb^{-1} , $\sqrt{(S+B)/S} = 1.4\%$
- The quest continues – newer det. simulation
 - use fancier generators: [TopRex](#), [Single Top](#)
 - Work-in-progress, unofficial results $\delta\sigma / \sigma \sim 10\%$








t-channel single top (ATLAS)

- ATLAS study of W-gluon single top (Dugan O'Neil's Ph.D. thesis):
 - ATLFAST parametrized detector simulation
 - Onetop+Pythia 5.72 signal, tt; M.E.+HERWIG Wbb, Wjj
 - Selections are rather similar to CMS study, plus:
 - $150 < M_{\text{top}} < 200 \text{ GeV}/c^2$
 - Result: $\sqrt{(S+B)/S} = 0.9\%$ with 10 fb^{-1} .
- What about V_{tb} ? Extracting $|V_{tb}|^2$ from x-section picks up uncertainties:
 - x-section uncertainty:
 - Statistical: 5% is a perhaps reasonable/optimistic guess for $L=10 \text{ fb}^{-1}$
 - Systematic $< 20\%$. Here things can easily be off by a lot, especially in the 1st year...
 - Luminosity uncertainty: traditionally 4-5%
 - Theoretical uncertainty:
 - From factorization and renormalization scale dependence (3% from NLO calculation)



s-channel single top (CMS)

- CMS study (S. Sablospitsky, CMS week 9/17/03)
 - disclaimer: work-in-progress, not yet 'official' results
 - use TopRex to generate single top, tt, Wbb. Pythia W+jets. Use CMSJET 4.801
 - selection:
 - one lepton with $E_T > 10$ GeV, $|\eta| < 2.5$
 - 2 B-tags with $E_T > 20$ GeV, no other jets with $E_T > 20$ GeV
 - Vector sum all final state P_T : $|\Sigma \vec{P}_T| < 15$ GeV
 - Reconst. top mass: $150 \text{ GeV} < M_{\text{top}} < 200 \text{ GeV}/c^2$
 - Results: $\sqrt{(S+B)/S} = 12\%$, $S/B = 8\%$ for $L = 30 \text{ fb}^{-1}$
 - which implies $\delta|V_{tb}| = 8.3\%$






s-channel single top (ATLAS)

- ATLAS study ([hep-ph/0003033](https://arxiv.org/abs/hep-ph/0003033))
 - Onetop+Pythia 5.72 signal, tt; M.E.+HERWIG Wbb, Wjj
 - Selection:
 - One isolated lepton with $P_T > 20$ GeV/c, $|\eta| < 2.5$
 - 2 b-jets with $E_T > 75$ GeV
 - Scalar sum all final state P_T : $|\Sigma P_T| < 175$ GeV
 - Reconst. top mass: $150 \text{ GeV} < M_{\text{top}} < 200 \text{ GeV}/c^2$
 - Results: $\sqrt{(S+B)/S} = 5.5\%$ for $L = 30 \text{ fb}^{-1}$, $\delta|V_{tb}| \sim 1\%$

- Wt single top production at ATLAS:
 - $65 < M_{jj} < 95 \text{ GeV}/c^2$
 - Event invariant mass $< 300 \text{ GeV}/c^2$
 - $\sqrt{(S+B)/S} = 4.4\%$ for $L = 30 \text{ fb}^{-1}$





Top Polarization in Single Top Events

- EW-produced top quarks are highly polarized:
 - In the top rest frame, its spin points along the direction of the down (d) quark.
 - Restrict to t-channel single top: 80% (69%) of the top (antitop) events have the 'd' quark in the final state. The angular distrib. of the charged lepton:
$$f(\cos \theta_\lambda) = \frac{1}{2} (1 + P \cos \theta_\lambda)$$
 - used signal samples with $P=+1$ and $P=-1$. Consider only W+jet background.
 - Chisquare fit, letting the two P contributions float.
 - The error on polarization measurement is 1.6% for 10 fb⁻¹ (Dugan's Ph.D. thesis)
- W boson helicity measurement:
 - Again restrict to t-channel signal.
 - Measure $\cos(\psi_l)$ distribution. ψ_l between the direction of the lepton in W rest frame and the direction of the W in the top rest frame.
 - Three component fit: f_L , f_R , f_{Long} , with $f_L + f_R + f_{\text{Long}} = 1$
 - Uncertainties of the order 2-3% for 30 fb⁻¹ of data.



Conclusions

- CDF and DØ accumulated more events than in entire Run I (factor of 2).
- Preliminary CDF study from 107 pb^{-1} sets promising single top limits
- However, many improvements needed if we are to observe single top in the next 2-3 years.
- Precision measurements of $|V_{tb}|$, as well as top polarization are not within (immediate) sight
- LHC will see very exciting results in the first (few) years of running:
 - $10 - 30 \text{ fb}^{-1}$ needed for observation in most searches.
 - Higher collider energy = considerable higher cross sections
- ATLAS and CMS single top studies are very mature
 - The Monte Carlo efforts are particularly strong: Top Rex, SingleTop.
- CDF/DØ can help with insight into systematics. Search strategies are already similar between Tevatron and CERN, but big improvements are just an idea away!